
Title	Effects of complex training on sprint, jump, and change of direction ability of soccer players: A systematic review and meta-analysis
Author(s)	Rohit K. Thapa, Danny Lum, Jason Moran and Rodrigo Ramirez-Campillo
Source	<i>Frontiers in Psychology, 11, Article 627869</i>

Copyright © 2021 Frontiers Media, Thapa, Lum, Moran and Ramirez-Campillo.



Effects of Complex Training on Sprint, Jump, and Change of Direction Ability of Soccer Players: A Systematic Review and Meta-Analysis

Rohit K. Thapa¹, Danny Lum^{2,3}, Jason Moran⁴ and Rodrigo Ramirez-Campillo^{5,6*}

¹ Department of Sports Biomechanics, Lakshmbai National Institute of Physical Education, Gwalior, India, ² Sport Science and Sport Medicine, Singapore Sport Institute, Singapore, Singapore, ³ Physical Education and Sport Science Academic Group, National Institute of Education, Nanyang Technological University, Singapore, Singapore, ⁴ School of Sport Rehabilitation and Exercises Sciences, University of Essex, Colchester, United Kingdom, ⁵ Human Performance Laboratory, Department of Physical Activity Sciences, Universidad de Los Lagos, Osorno, Chile, ⁶ Centro de Investigación en Fisiología del Ejercicio, Facultad de Ciencias, Universidad Mayor, Santiago, Chile

OPEN ACCESS

Edited by:

Miguel-Angel Gomez-Ruano,
Polytechnic University of
Madrid, Spain

Reviewed by:

Souhail Hermassi,
Qatar University, Qatar
Tomás T. Freitas,
Catholic University San Antonio of
Murcia, Spain

*Correspondence:

Rodrigo Ramirez-Campillo
r.ramirez@ulagos.cl
orcid.org/0000-0003-2035-3279

Specialty section:

This article was submitted to
Movement Science and Sport
Psychology,
a section of the journal
Frontiers in Psychology

Received: 10 November 2020

Accepted: 24 December 2020

Published: 22 January 2021

Citation:

Thapa RK, Lum D, Moran J and
Ramirez-Campillo R (2021) Effects of
Complex Training on Sprint, Jump,
and Change of Direction Ability of
Soccer Players: A Systematic Review
and Meta-Analysis.
Front. Psychol. 11:627869.
doi: 10.3389/fpsyg.2020.627869

The aim of this meta-analysis was to evaluate the effects of complex training (CT) on sprint, jump, and change of direction (COD) ability among soccer players. After an electronic search, 10 peer-reviewed articles were considered in the meta-analysis. The athletes included in this meta-analysis were amateur to professional level male soccer players (age range, 14–23 years). These studies incorporated CT in soccer players who were compared to a control group. Significant moderate to large improvements were observed in the CT group [sprint: standard mean difference (SMD) = 0.92–1.91; jump: SMD = 0.96–1.58; COD: SMD = 0.97–1.49] when compared to control groups. Subgroup analysis were also conducted based on age, duration, and competitive level. The beneficial effects of CT were greater in players <18 vs. ≥18 years (linear sprinting; SMD = 2.01 vs. –0.13), after ≥8 vs. <8 weeks (jumping and COD; SMD = 1.55–2.01 vs. 0.31–0.64, respectively) and among professional vs. amateur players (linear sprinting and with COD; SMD = 1.53–1.58 vs. 0.08–0.63, respectively). In conclusion, regular soccer training programs may be supplemented with CT to improve sprint, jump, and COD performance. A longer duration of CT (≥8 weeks) seems to be optimal in improving the physical abilities of soccer players. Professional players and <18 years players may benefit more from CT program.

Keywords: football, plyometric exercise, post-activation performance enhancement, physical education and training, resistance training, sports

INTRODUCTION

Soccer is a team sport which requires players to execute several short-duration maximal- and near-maximal physical efforts such as sprinting, jumping, and change(ing) of direction (COD) in order to overcome opponents during play (Stølen et al., 2005; Barnes et al., 2014). These physical fitness abilities are crucial in determining a player's performance potential during soccer games and tournaments (Castagna et al., 2003; Arnason et al., 2004; Faude et al., 2012), and have been

linked with strength and power of the lower extremities (Arnason et al., 2004; Wisløff et al., 2004). To increase strength and power, resistance training (RT) and plyometric (PT)/power training programs are widely used in soccer (De Hoyo et al., 2016; Bauer et al., 2019; Ramirez-Campillo et al., 2020b). The aforementioned training methods induce neuromuscular adaptations, such as enhanced stretch-shortening cycle function, motor unit recruitment, firing frequency, intra- and inter-muscular coordination, and morphological changes (e.g., fiber type or pennation angle), enhancing overall performance (Markovic and Mikulic, 2010; Cormie et al., 2011). Although both RT (Marios et al., 2006; Sander et al., 2013; Silva et al., 2015) and PT (Van De Hoef et al., 2019; Ramirez-Campillo et al., 2020a,b) may improve strength and power, respectively, conducting independent training sessions for each method within a congested weekly micro-cycle may be troublesome for players and coaches.

Alternatively, a pragmatic approach combines both RT and PT methods within a single training work set, allowing more time to be devoted to specific soccer training activities. One such combination is termed as complex training (CT) (Fleck and Kontor, 1986), which commonly involves the performance of an exercise set with a high-load RT exercise, followed immediately by the execution of a low-load plyometric-power exercise. This format of training usually involves the combination of two biomechanically similar exercises with the high-load resistance exercise performed first [e.g., squat at 90% of one repetition maximum (1RM)], followed by the low-load power exercise (e.g., squat jumps) (Fleck and Kontor, 1986; Docherty et al., 2004). Such exercise sequencing stimulates the post-activation potentiation of performance, a phenomenon (Docherty et al., 2004; Hodgson et al., 2005; Carter and Greenwood, 2014; Prieske et al., 2020), that stimulates motor unit recruitment thus increasing the force producing potential of the utilized musculature within a given movement (Healy and Comyns, 2017).

In the last four decades (Fleck and Kontor, 1986), a considerable amount of published studies have analyzed the effects of CT on athletes' physical fitness and some meta-analyses have attempted to congregate the extant literature (Freitas et al., 2017; Bauer et al., 2019; Cormier et al., 2020; Pagaduan and Pojskic, 2020). Some meta-analyses have also been conducted specifically on team sports athletes (Freitas et al., 2017; Cormier et al., 2020). However, these studies included athletes with different sporting backgrounds (i.e., soccer, futsal, basketball, volleyball, rugby, football, water polo, track and field, and handball), potentially biasing the results for a particular sport, such as soccer. Indeed, the effects of PT and RT may vary according to an athlete's athletic history (Izquierdo et al., 2002; Fleck and Kraemer, 2004; De Villarreal et al., 2012; Asadi et al., 2016; Taylor et al., 2018; Vlachopoulos et al., 2018). Moreover, a 5-fold difference may be observed in the magnitude of the effects of CT on the physical fitness performance (i.e., sprinting) of athletes with different team sport background (Freitas et al., 2017).

Therefore, in order to clarify the specific effects of CT (and its moderators; e.g., programme duration) on soccer player's

physical fitness performance, the aim of this meta-analysis was to evaluate the effects of CT interventions on sprint, jump, and COD ability among soccer players. With reference to previous studies (Freitas et al., 2017; Kobal et al., 2017; Bauer et al., 2019; Cormier et al., 2020), we hypothesized that CT would be effective in improving soccer players' physical fitness, with greater effects observed as compared to a control condition. Accordingly, our research question was: what are the effects of CT interventions (and its moderators) on sprint, jump, and COD ability among soccer players when compared to a control group?

METHODS

The lead investigator (RKT) and a research assistant conducted electronic searches. The PubMed and Google scholar databases were used, conforming to the guidelines set by the Preferred Reported Items for Systematic Reviews and Meta-Analysis (PRISMA) (Moher et al., 2015). This study was registered with the International Platform of Registered Systematic Review and Meta-Analysis Protocols (INPLASY202090079). Articles published up to August 30th, 2020 were considered. Keywords were selected through experts' opinion and a systematic literature review. Using Boolean logic, the following combination of keywords was used in the search databases: "complex training" or "contrast training" or "combination of strength training and plyometrics" and "soccer" or "football." An example for search strategy used in PubMed was: [(((complex training) OR contrast training) OR (combination of strength training and plyometrics)) AND soccer] OR football. Relevant articles' reference lists were examined to identify further articles for inclusion in the meta-analysis.

Inclusion Criteria and Exclusion Criteria

A PICOS (participants, intervention, comparators, outcomes, and study design) approach was used to rate studies' eligibility (Liberati et al., 2009). The respective inclusion/exclusion criteria adopted in our meta-analysis are reported in **Table 1**.

Methodological Quality of Studies

The Tool for the Assessment of Study Quality and Reporting in Exercise (TESTEX) was used to assess the quality of studies included in this meta-analysis (Smart et al., 2015). This tool comprises of 12 methodological quality assessment criteria (a full description of each criteria is provided in the results section for the included studies). The maximum score a study can obtain is 15 points, with higher ratings reflecting better study quality and reporting (Smart et al., 2015).

Data Extraction

We extracted from each eligible study, data relating to linear sprints over various distances (5–40-m linear sprint tests), vertical jumping and sprints which incorporated a COD (505, S4×5, T-test, BAT, S180 tests and dribbling with or without ball) (Castagna et al., 2003; Arnason et al., 2004; Faude et al., 2012). Means, standard deviations (SD), and sample sizes (*n*) (**Table 2**) were extracted by one author (RKT) from the included papers and were corroborated by a second author (DL). Any discrepancy

TABLE 1 | Selection criteria used in the meta-analysis.

Category	Inclusion criteria	Exclusion criteria
Population	Apparently healthy soccer players, with no restrictions on their playing level, sex, or age	Soccer players with health problems (e.g., injuries, recent surgery)
Intervention	A complex training programme, defined as a combination of heavy load strength exercise followed by a low load plyometric/power exercise, set by set	Exercise interventions not involving complex training or exercise interventions involving contrast training, where strength training exercises were conducted first and plyometric/power exercises at the end of the session (or during a different session)
Comparator	Active control group	Absence of active control group
Outcome	At least one measure of physical fitness (linear sprinting, jumping, and change of direction speed) before and after the training intervention	Lack of baseline and/or follow-up data
Study design	Controlled trials	Non-controlled trials

between the authors was resolved through discussion with a third author (RRC). Where data were displayed in a figure or no numerical data were provided by authors after being contacted, validated ($r = 0.99$, $p < 0.001$) software (WebPlotDigitizer; <https://apps.automeris.io/wpd/>) was used to derive numerical data from figures (Drevon et al., 2017). In addition to the study data, sample characteristics, including age, playing level, training frequency, total duration of the intervention, and the type of training protocol used in the study were extracted and recorded.

Statistical Analysis

The meta-analysis was conducted using the Review Manager Software (RevMan 5.3). Statistical significance was set at $p \leq 0.05$. The inverse-variance random effects model for meta-analyses was used because it allocates a proportionate weight to trials based on the size of their individual standard errors (Deeks et al., 2008) and facilitates analysis whilst accounting for heterogeneity across studies (Kontopantelis et al., 2013). Effect sizes are represented by the standardized mean difference and are presented alongside 95% confidence intervals (CI). Standardized mean differences were calculated using the following equation (Cohen, 1988):

$$\text{SMD} = \frac{M1 - M2}{\text{SD pooled}}$$

Where $M1 - M2$ is the difference in the mean outcome between groups, and SD pooled is the pooled standard deviation of the outcome among subjects, which was calculated using the following equation:

$$\text{SD pooled} = \sqrt{\frac{SD_1^2 + SD_2^2}{2}}$$

Where SD_1^2 is the squared standard deviation of group 1 and SD_2^2 is the squared standard deviation of group 2.

The SMD were interpreted with threshold values as follow: <0.2 , trivial; $0.2-0.6$, small; $>0.6-1.2$, moderate; $>1.2-2.0$, large; $>2.0-4.0$, very large; >4.0 , extremely large (Hopkins et al., 2009).

In studies with more than one intervention group, the control group was proportionately divided to facilitate comparison across all participants (Higgins et al., 2008).

Subgroup analyses were performed on the variables that could have influenced the outcome after CT interventions, with median values of continuous variables used as cut-off values for grouping [age (<18 vs. ≥ 18 years), and CT programme duration (<8 vs. ≥ 8 weeks)]. The subgrouping of studies according to the athlete's competition level [professional vs. amateur (youth, university); as reported in the studies] was also considered. The analyses were conducted when there were at least one study in the subgroups.

The I^2 statistics was used to assess the heterogeneity among the studies, with values of <25 , $25-75$, and $>75\%$ interpreted as low, moderate and high heterogeneity, respectively (Higgins et al., 2003; Higgins, 2008). Heterogeneity was assessed in line with a low chi-squares' p -value (Higgins et al., 2003; Higgins, 2008) and a high Tau² value (Borenstein et al., 2009).

RESULTS

The initial search resulted in the retrieval of 1,373 articles and 22 additional articles were extracted through other sources. After removing duplicates, meta-analysis and systematic reviews 547 articles remained. After screening titles and abstracts for relevance, 516 articles were excluded with 31 full texts being retained. Further screening of articles based on our inclusion and exclusion criteria resulted in the inclusion of 10 studies in the meta-analysis (Maio Alves et al., 2010; Faude et al., 2013; Brito et al., 2014; Cavaco et al., 2014; García-Pinillos et al., 2014; Hammami et al., 2017a,b, 2019; Chatzinikolaou et al., 2018; Ali et al., 2019) (Figure 1).


General Characteristics of Studies

The study characteristics are presented in Table 3. A total of 313 subjects (all males) were included in this meta-analysis with 188 professional players and 125 amateur players. Sprinting time was measured through 5–40-m linear sprint tests. Jumping was measured using jump height in the squat jump and countermovement jump. The sprint time with COD was measured among the included studies using the 505, S4×5, T-test, BAT, and S180 tests, with or without dribbling a soccer ball. Considering the different number of COD (i.e., turns) involved among the tests, in a *posteriori* decision the tests were further divided based on those including ≤ 3 turns and > 3 turns.

TABLE 2 | The mean \pm standard deviation of fitness variables reported for the complex training and control conditions in the included studies.

References	Fitness attribute	Complex training			Control		
		Pre	Post	n	Pre	Post	n
Ali et al. (2019)	20m (s)	3.48 \pm 0.35	3.23 \pm 0.26	12	3.40 \pm 0.26	3.54 \pm 0.27	12
	CMJ (cm)	46.98 \pm 4.84	50.75 \pm 4.91	12	43.97 \pm 6.92	44.35 \pm 6.92	12
	T-test (s)	11.72 \pm 1.11	11.15 \pm 0.94	12	11.60 \pm 1.01	11.73 \pm 1.16	12
Maio Alves et al. (2010) (1 session/week)	5m (s)	1.09 \pm 0.07	0.99 \pm 0.03	9	1.13 \pm 0.08	1.11 \pm 0.03	6
	15m (s)	2.56 \pm 0.10	2.38 \pm 0.09	9	2.59 \pm 0.07	2.56 \pm 0.02	6
	SJ (cm)	41.02 \pm 6.11	46.19 \pm 7.70	9	41 \pm 3.05	40.7 \pm 3.91	6
	CMJ (cm)	42.84 \pm 4.55	42.92 \pm 5.56	9	42.63 \pm 3.35	41.53 \pm 2.71	6
	505 (s)	2.34 \pm 0.11	2.31 \pm 0.09	9	2.37 \pm 0.09	2.39 \pm 0.15	6
Maio Alves et al. (2010) (2 sessions/week)	5m (s)	1.13 \pm 0.04	1.06 \pm 0.03	8	1.13 \pm 0.08	1.11 \pm 0.03	6
	15m (s)	2.57 \pm 0.10	2.49 \pm 0.07	8	2.59 \pm 0.07	2.56 \pm 0.02	6
	SJ (cm)	39.68 \pm 4.34	43.5 \pm 4.50	8	41 \pm 3.05	40.7 \pm 3.91	6
	CMJ (cm)	41.78 \pm 5.28	42.79 \pm 4.45	8	42.63 \pm 3.35	41.53 \pm 2.71	6
	505 (s)	2.32 \pm 0.08	2.32 \pm 0.03	8	2.37 \pm 0.09	2.39 \pm 0.15	6
Brito et al. (2014)	5m (s)	1.13 \pm 0.02	1.02 \pm 0.02	12	1.14 \pm 0.02	1.11 \pm 0.02	26
	20m (s)	3.25 \pm 0.09	3.05 \pm 0.07	12	3.21 \pm 0.13	3.17 \pm 0.12	26
Cavaco et al. (2014) (1 session/week)	15m (s)	2.72 \pm 0.25	2.59 \pm 0.18	5	2.68 \pm 0.19	2.66 \pm 0.18	6
	Agility with ball (s)	10.64 \pm 1.82	9.80 \pm 1.43	5	9.88 \pm 0.48	9.90 \pm 0.46	6
Cavaco et al. (2014) (2 sessions/week)	15m (s)	2.6 \pm 0.14	2.46 \pm 0.15	5	2.68 \pm 0.19	2.66 \pm 0.18	6
	Agility with ball (s)	9.64 \pm 1.23	8.54 \pm 0.37	5	9.88 \pm 0.48	9.90 \pm 0.46	6
Chatzinikolaou et al. (2018)	10m (s)	1.90 \pm 0.08	1.83 \pm 0.07	12	1.93 \pm 0.12	2.02 \pm 0.12	10
	30m (s)	4.8 \pm 0.43	4.32 \pm 0.21	12	4.69 \pm 0.43	4.79 \pm 0.44	10
	CMJ (cm)	35.84 \pm 4	38.2 \pm 3.84	12	35.03 \pm 3.55	32.08 \pm 2.67	10
	CMJ arm swing (cm)	42.33 \pm 4.74	45.4 \pm 4.37	12	41.49 \pm 3.07	38.33 \pm 1.76	10
	SJ (cm)	32.95 \pm 4.19	34.95 \pm 3.81	12	32.29 \pm 3.9	30.1 \pm 3.61	10
Faude et al. (2013)	10m (s)	1.77 \pm 0.07	1.77 \pm 0.05	8	1.73 \pm 0.08	1.77 \pm 0.05	8
	30m (s)	4.26 \pm 0.15	4.27 \pm 0.09	8	4.12 \pm 0.10	4.19 \pm 0.09	8
	CMJ (cm)	40.2 \pm 4.8	41.4 \pm 3.5	8	43 \pm 2.1	39.8 \pm 2.4	8
	COD (s)	9.8 \pm 0.7	9.4 \pm 0.4	8	9.5 \pm 0.04	9.3 \pm 0.3	8
García-Pinillos et al. (2014)	5m (s)	1.67 \pm 0.23	1.42 \pm 0.18	17	1.66 \pm 0.17	1.46 \pm 0.18	13
	10m (s)	2.47 \pm 0.25	2.14 \pm 0.30	17	2.42 \pm 0.15	2.24 \pm 0.17	13
	20m (s)	3.83 \pm 0.29	3.51 \pm 0.24	17	3.74 \pm 0.17	3.53 \pm 0.21	13
	30m (s)	5.11 \pm 0.35	4.79 \pm 0.30	17	4.96 \pm 0.18	4.77 \pm 0.24	13
	CMJ (cm)	42 \pm 6	45 \pm 4	17	45 \pm 3	46 \pm 3	13
	BAT (s)	12.29 \pm 0.54	11.66 \pm 0.43	17	11.93 \pm 0.58	11.89 \pm 0.46	13
Hammami et al. (2017a)	5m (s)	1.14 \pm 0.03	1.01 \pm 0.02	16	1.15 \pm 0.09	1.12 \pm 0.06	12
	10m (s)	1.87 \pm 0.23	1.73 \pm 0.13	16	1.92 \pm 0.13	1.90 \pm 0.14	12
	20m (s)	3.32 \pm 0.17	3.04 \pm 0.04	16	3.30 \pm 0.22	3.30 \pm 0.23	12
	30m (s)	4.57 \pm 0.14	4.23 \pm 0.06	16	4.59 \pm 0.33	4.56 \pm 0.33	12
	40m (s)	5.96 \pm 0.21	5.42 \pm 0.15	16	5.94 \pm 0.41	5.85 \pm 0.42	12
	SJ (cm)	36.8 \pm 2.6	45.4 \pm 3.5	16	35.4 \pm 5.1	34.3 \pm 5.2	12
	CMJ (cm)	38.4 \pm 1.9	48.1 \pm 4.5	16	3.7 \pm 6.8	36.1 \pm 5.9	12
	S180 (s)	8.37 \pm 0.21	7.93 \pm 0.26	16	8.41 \pm 0.33	8.40 \pm 0.35	12
	S4x5 (s)	6.28 \pm 0.25	5.83 \pm 0.04	16	6.29 \pm 0.21	6.26 \pm 0.16	12
Hammami et al. (2017b)	5m (s)	1.14 \pm 0.04	1.01 \pm 0.02	17	1.06 \pm 0.12	1.11 \pm 0.11	12
	10m (s)	1.91 \pm 0.4	1.76 \pm 0.04	17	1.87 \pm 0.16	1.90 \pm 0.17	12
	20m (s)	3.30 \pm 0.17	3.04 \pm 0.05	17	3.21 \pm 0.25	3.27 \pm 0.25	12
	30m (s)	4.56 \pm 0.14	4.23 \pm 0.08	17	4.51 \pm 0.37	4.56 \pm 0.37	12
	40m (s)	5.93 \pm 0.21	5.43 \pm 0.16	17	5.90 \pm 0.40	5.81 \pm 0.43	12
	S180 (s)	8.37 \pm 0.21	7.94 \pm 0.25	17	8.39 \pm 0.35	8.40 \pm 0.37	12
	S4x5 (s)	6.24 \pm 0.26	5.84 \pm 0.05	17	6.18 \pm 0.29	6.13 \pm 0.22	12
Hammami et al. (2019)	5m (s)	1.13 \pm 0.04	1.01 \pm 0.17	14	1.03 \pm 0.06	1.08 \pm 0.04	12
	40m (s)	5.93 \pm 0.22	5.43 \pm 0.17	14	5.85 \pm 0.32	5.75 \pm 0.33	12
	SJ (cm)	36.8 \pm 2.8	45.4 \pm 3.7	14	36.2 \pm 3.8	34.8 \pm 4.5	12
	CMJ (cm)	38.2 \pm 2.1	47 \pm 5.9	14	37.9 \pm 5.4	37.2 \pm 4.5	12
	S4x5 (s)	6.25 \pm 0.26	5.83 \pm 0.47	14	6.15 \pm 0.25	6.61 \pm 0.26	12

5m, 5-m linear sprint time; 10m, 10-m linear sprint time; 20m, 20-m linear sprint time; 30m, 30-m linear sprint time; 40m, 40-m linear sprint time; BAT, Balsa agility test; COD, change of direction; CMJ, countermovement jump; CMJ arm swing, countermovement jump with arm swing; SJ, squat jump; S180, Sprinting 9-3-6-3-9m with 180° Turns; S4x5, 4x5m sprints with COD.



Accordingly, the 505 and S4×5 tests involved ≤ 3 turns, while the T-test, BAT, and S180 tests involved > 3 turns. Two studies assessed COD with ball, using ≤ 3 turns (Faude et al., 2013) and > 3 turns (Cavaco et al., 2014).

Methodological Quality of Included Studies

Table 4 shows the methodological quality of the eligible studies in this meta-analysis. The score of all the studies according to TESTEX criteria ranged from 10 to 13 points. All studies selected for the meta-analysis obtained a score ≥ 10 points. Therefore, no study was excluded based on its methodological quality.

Meta-Analysis Results for Linear Sprint Performance

A moderate to very large significant improvement was noted after CT (within group, pre-post analysis) for 5-m (SMD = 2.44, 95% CI = 1.34–3.54, $p < 0.001$), 10-m (SMD = 0.67, 95% CI = 0.34–0.99, $p < 0.001$), 15-m (SMD = 1.05, 95% CI = 0.46–1.64, $p < 0.001$), 20-m (SMD = 1.43, 95% CI = 0.74–2.12, $p < 0.001$), 30-m (SMD = 1.31, 95% CI = 0.27–2.35, $p = 0.01$), and 40-m linear sprint performance (SMD = 2.65, 95% CI = 2.08–3.23, $p < 0.001$). The relative weight of each study in the analyses varied

TABLE 3 | Study characteristics.

References	Groups	Subjects (n)	Age (Mean ± SD)	Training intervention			Variables	
				Level	Frequency/ wk.	Duration (weeks)		
Maio Alves et al. (2010)	CT1 CT2 CG	9 8 6	17.4 ± 0.6	P	1 2	6	Pre-season 1st station: 6 rep of 90 squat exercise at 85% 1RM + 1 set of 5 m high skipping in straight line and then 5 m sprint. 2nd station: 6 rep of calf extension exercise at 90% 1RM + 8 vertical jumps + 3 high-ball headers. 3rd station: 6 rep of leg extension exercise at 80% 1RM + 6 jumps from the seated position + 3 drops jumps (60 cm) executing a soccer heading. The load was increased 5% (from 1RM) every 2 weeks.	5m, 15m, SJ, CMJ, COD
Ali et al. (2019)	CT CST CG	12 12 12	22 ± 2.4 20.8 ± 2.1 21.5 ± 1.8	C	3	6	Off-season Squats (80% 1 RM) 3 × 12 + Depth jumps 3 × 12 Barbell lunge (80% 1 RM) 3 × 12 + Split squat jumps 3 × 12 Lateral lunge (80% 1 RM) 3 × 12 + Lateral hops 3 × 12	20m, CMJ, COD (T-test)
Brito et al. (2014)	CG ST PT CT	21 12 12 12	20.3 ± 1.6	C	2	9	In-season Station 1: 6 reps of 85% of 1RM SQ at 90° + 1 set high skipping, cyclically, with thighs Parallel to the ground (5m) + 1 5m sprint Station 2: 6 reps at 90% of 1RM calf extension + 8 consecutive vertical jumps with Minimum GCT + 3 ball headers (jumping as high as possible) Station 3: 6 reps at 80% of 1RM of leg extension + 6 vertical jumps from seated position + 3 DJ (60 cm) with soccer heading post vertical jumps	5m, 20m, SJ, CMJ, COD (T-test)
Cavaco et al. (2014)	CT1 CT2 CG	5 5 6	13.8 ± 0.4 14.2 ± 0.4 14.2 ± 0.8	Y	1 2	6	First station: 3 sets, with 6 squat repetitions at 85% of 1-RM, 15 m plus top speed and cross, separated by 3 min of rest. Second station: 3 sets, 6 repetitions of the squat at 85% of 1-RM, plus agility with the ball and shot at the goal, separated by 3 min of rest. The load in the squat exercise was increased by 5% from initial 1RM each 2 weeks.	15m, 15m agility with ball
Chatzinikolaou et al. (2018)	CG CT	10 12	14.1 ± 0.6 14.3 ± 0.7	P	2	5	Off-season 2 days ST with 13 exercises 2 days CXT Day 1: Barbell cleans+10–20m sprints/agility drills; Kettlebell snatch+10–20m sprints/agility drills; Box jumps+10m agility drills Day 2: Power bag jerk+10–20m sprints; Kettlebell snatch_10–20m sprints; Resistive sprints + sprints (8m + 15–20m)	10m, 30m, SJ, CMJ, COD (Arrowhead)
Faude et al. (2013)	CG CT	8 8	22.5 ± 2.5	A	2	7	In-season Day 1: Unilateral load half SQ (90% of estimated 1 RM) + 5 single leg hurdle jumps (right and left leg) Day 2: (50–60% 1RM)	10m, 30m, CMJ, COD

(Continued)

TABLE 3 | Continued

References	Groups	Subjects (n)	Age (Mean ± SD)	Training intervention			Variables	
				Level	Frequency/ wk.	Duration (weeks)		
García-Pinillos et al. (2014)	CG	13	15.9 ± 1.4	P	2	12	5 loaded half SQ + 5 DJ + 5m sprint; 5 loaded calf raises + 5 high straight jumps + 1 headers; 8 loaded lateral half SQ (4 left, 4 right side) + 8 lateral jumps + 10 zig-zag sprints; 8 loaded step ups (4 left, 4 right side) + 4 bounding jumps + 3 headers In-season Weeks 1–2: 90° isometric half SQ (40 s) + jump from seated position (6 reps) Weeks 3–4: 90° isometric half SQ (60 s) + alternate single leg jump (6 reps) Weeks 5–6: 90° isometric half SQ (80 s) + jump from seated position/ alternate single leg jump (alternatively) Weeks 7–8: 90° isometric half SQ (40 s) + alternate single leg jump (6 reps) + 90° isometric half SQ (40 s) + jump from seated position (6 reps) Week 9–10: 90° isometric half SQ (40 s) + alternate single leg jump (6 reps) + 90° isometric half SQ (40 s) + jump from seated position (6 reps) Week 11–12: same as 9–10	5m, 10m, 20m, 30m, CMJ, COD (BAT)
	CT	17						
Hammami et al. (2017a)	CG	12	16 ± 0.5	P	2	8	In-season First 4 weeks: Back Half SQ+3 consecutive CMJ Second 4 weeks: Back Half SQ+1 CMJ+15m sprint	5m, 10m, 20m, 30m, 40m, SJ, CMJ, S180, S4x5m
	ST	16						
	CT	16						
Hammami et al. (2017b)	CT	17	16.0 ± 0.5	P	2	8	In-season Weeks 1–4: Half SQ + 3 consecutive CMJ Weeks 5–8: Half SQ + 1 CMJ + 15m sprint	5m, 10m, 20m, 30m, 40m, S4x5, S180
	CG	12						
Hammami et al. (2019)	CT	14	15.8 ± 0.4	P	2	8	In-season Weeks 1–4: Half SQ + 3 consecutive CMJ Weeks 5–8: Half SQ + 1 CMJ + 15m sprint	5m, 40m, SJ, CMJ, COD (4x5m)
	PT	14						
	CG	12						

1RM, one repetition maximum; 5m, 5-m linear sprint time; 10m, 10-m linear sprint time; 15m, 15-m linear sprint time; 20m, 20-m linear sprint time; 30m, 30-m linear sprint time; 40m, 40-m linear sprint time; A, amateur soccer team; BAT, Balsom agility test; C, college level; CG, control Group; CMJ, countermovement jump; COD, change of direction; CT, complex training group; CT1, complex training with one session per week; CT2, complex training with two sessions per week; CST, contrast training group; DJ, drop jump; P, professional soccer team. Some youth teams were categorized as professionals, considering the training load and/or playing level described in the respective studies (e.g., "Elite"); PT, plyometric training group; RDL, Romanian deadlift; rep, repetitions; S, semi-professional; S180, Sprinting 9-3-6-3-9m with 180° Turns; S4x5, 4x5m sprints with COD; SJ, squat jump; SPT, sprint training group; SQ, squat; Y, youth.

between 10.8 and 37.7%, with low to high heterogeneity ($I^2 = 0-88.0\%$).

A moderate to large significant difference was noted between CT and control (between group, post analysis) for 5-m (SMD = 1.91, 95% CI = 0.81–3.00, $p < 0.001$, **Figure 2A**), 10-m (SMD = 0.92, 95% CI = 0.32–1.53, $p = 0.003$, **Figure 2B**), 15-m (SMD = 1.07, 95% CI = 0.31–1.83, $p = 0.006$, **Figure 2C**), 20-m (SMD = 1.04, 95% CI = 0.49–1.58, $p < 0.001$, **Figure 2D**), and 40-m linear sprint performance (SMD = 1.28, 95% CI = 0.80–1.76, $p < 0.001$, **Figure 2F**). The relative weight of each study in the analyses varied between 10.1 and 35.3%, with low to high heterogeneity ($I^2 = 0-87\%$). Only 30-m linear sprint showed no difference between CT and control (**Figure 2E**).

Meta-Analysis Results for Vertical Jump Performance

A moderate to large significant improvement was noted after CT (within group, pre-post analysis) for countermovement jump (SMD = 0.89, 95% CI = 1.43–0.35, $p < 0.001$) and squat jump height performance (SMD = 1.33, 95% CI = 2.12–0.54, $p = 0.001$). The relative weight of each study in the analyses varied between 10.4 and 17.9%, with low to high heterogeneity ($I^2 = 64.0-76.0\%$).

In addition, a moderate to large significant difference was noted between CT and control (between group, post analysis) for countermovement jump (SMD = 0.96, 95% CI = 1.64–0.29, $p = 0.005$, **Figure 3A**) and squat jump height performance (SMD =

TABLE 4 | Methodological quality score of the studies included in the meta-analysis.

References	Items												Total points (from a maximum of 15)	
	1	2	3	4	5	6	7	8	9	10	11	12		
Ali et al. (2019)	1	–	–	1	–	1	1	2	1	1	1	1	1	10
Maio Alves et al. (2010)	1	–	1	1	–	1	1	2	1	1	1	1	1	11
Brito et al. (2014)	1	1	1	1	–	1	1	2	1	1	1	1	1	12
Cavaco et al. (2014)	1	1	1	1	–	1	1	2	1	1	1	1	1	12
Chatzinikolaou et al. (2018)	1	1	1	1	–	2	1	2	1	1	1	1	1	13
Faude et al. (2013)	1	1	1	–	–	2	1	2	1	1	1	1	1	12
García-Pinillos et al. (2014)	1	1	1	1	–	1	1	2	1	1	1	1	1	12
Hammami et al. (2017a)	1	1	1	–	–	1	1	2	1	1	1	1	1	11
Hammami et al. (2017b)	1	1	1	1	–	1	1	2	1	1	1	1	1	11
Hammami et al. (2019)	1	1	1	1	–	1	1	2	1	1	1	1	1	12

Note for items 1 through 12: (1) eligibility criteria specified (1 point); (2) randomization defined (1 point); (3) allocation concealment (1 point); (4) groups similar at baseline (1 point); (5) assessor blinding in study reporting (1 point); (6) outcome measures assessed in 85% of patients (3 points); (7) intention-to-treat analysis (1 point); (8) between-group statistical comparisons reported (2 points); (9) point measures and measures of variability for all reported outcome measures (1 point); (10) activity monitoring in the controlled group (1 point); (11) relative exercise intensity remained constant (1 point); (12) exercise volume and energy expenditure (1 point).




FIGURE 2 | Forest plot of standardized mean difference in post-intervention outcomes between complex training group and control group, including (A) 5-m sprint time, (B) 10-m sprint time, (C) 15-m sprint time, (D) 20-m sprint time, (E) 30-m sprint time, (F) 40-m sprint time. Note: the relative weight of each study in the analysis is indicated by the size of the green squares.

1.58, 95% CI = 2.38–0.78, $p < 0.001$, **Figure 3B**). The relative weight of each study in the analyses varied between 10.4 and 23.1%, with moderate heterogeneity ($I^2 = 60.0\text{--}73.0\%$).

Meta-Analysis Results for Change of Direction Performance

A moderate to large significant improvement was noted after CT (within group, pre-post analysis) for COD with ≤ 3 turns (SMD = 1.11, 95% CI = 0.33–1.88, $p = 0.005$) and COD with > 3 turns (SMD = 1.24, 95% CI = 0.83–1.64, $p < 0.001$). The relative weight of each study in the analyses varied between 8.7 and 19.5%, with moderate to high heterogeneity ($I^2 = 26.0\text{--}77.0\%$).

In addition, a moderate to large significant difference was noted between CT and control (between group, post analysis) for COD with ≤ 3 turns (SMD = 1.49, 95% CI = 0.40–2.58, $p = 0.007$, **Figure 4A**) and COD with > 3 turns (SMD = 0.97, 95% CI = 0.41–1.53, $p < 0.001$, **Figure 4B**). The relative weight of each study in the analysis varied between 4.5 and 23.0%, with moderate to high heterogeneity ($I^2 = 47.0\text{--}82.0\%$).

Subgroup Analyses

A subgroup analysis was performed based on age (< 18 vs. ≥ 18 years), playing level (professional vs. amateur), and duration




FIGURE 3 | Forest plot of standardized mean difference in post-intervention outcomes between complex training group and control group, including **(A)** countermovement jump (CMJ) height, **(B)** squat jump (SJ) height. Note: the relative weight of each study in the analyses is indicated by the size of the green squares.




FIGURE 4 | Forest plot of standardized mean difference in post-intervention outcomes between complex training group and control group, including **(A)** change of direction time with ≤3 turns, and **(B)** change of direction time with >3 turns. Note: the relative weight of each study in the analyses is indicated by the size of the green squares.

of the training intervention (<8 vs. ≥8 weeks) as covariates (Table 5).

Regarding the chronological age of players, a significant difference ($p < 0.001$, Table 5) in 30-m linear sprint time was

noted between players ≥18 years (SMD = -0.13) and <18 years (SMD = 2.01), in favor of younger players, after CT.

For training programme duration, a greater improvement in countermovement jump (SMD = 1.54 vs. 0.40, $p = 0.02$), squat

TABLE 5 | Subgroup analyses of potential moderator factors for linear sprint time, vertical jump height and sprint with change of direction time.

	Groups (n)	SMD (95% CI)	I ²	p	p _{diff}
5 m sprint					
Age					
≥18	2	2.84 (−1.85–7.54)	95	0.24	0.85
<18	6	2.37 (1.21–3.53)	87	<0.001	
Duration					
≥8 weeks	6	2.69 (1.22–4.15)	92	<0.001	0.31
<8 weeks	2	1.82 (0.98–2.65)	0	<0.001	
10 m sprint					
Age					
≥18	2	0.21 (−0.45–0.87)	0	0.53	0.12
<18	4	0.81 (0.44–1.18)	0	<0.001	
Playing level					
Professional	5	0.75 (0.41–1.09)	0	<0.001	0.16
Amateur	1	0.00 (−0.98–0.98)	–	1.00	
Duration					
≥8 weeks	4	0.72 (0.34–1.09)	0	<0.001	0.63
<8 weeks	2	0.48 (−0.39–1.36)	46	0.28	
20 m sprint					
Age					
≥18	3	1.06 (−0.15–2.28)	80	0.09	0.32
<18	3	1.76 (1.11–2.42)	47	<0.001	
Playing level					
Professional	4	1.38 (0.49–2.27)	78	0.002	0.86
Amateur	2	1.55 (−0.03–3.13)	81	0.05	
Duration					
≥8 weeks	5	1.56 (0.76–2.36)	76	<0.001	0.19
<8 weeks	1	0.78 (−0.05–1.62)	–	0.07	
30 m sprint					
Age					
≥18	2	−0.13 (−0.79–0.52)	0	0.69	<0.001
<18	4	2.01 (0.97–3.06)	81	<0.001	
Playing level					
Professional	5	1.58 (0.46–2.71)	87	0.006	0.03
Amateur	1	−0.08 (−1.06–0.90)	–	0.88	
Duration					
≥8 weeks	4	1.64 (0.19–3.10)	90	0.03	0.34
<8 weeks	2	0.66 (−0.76–2.08)	78	0.36	
CMJ					
Age					
≥18	3	−0.71 (−1.43 to −0.35)	0	<0.001	0.57
<18	6	−0.99 (−1.79 to −0.19)	80	0.02	
Playing level					
Professional	7	−1.00 (−1.68 to −0.32)	76	<0.001	0.35
Amateur	2	−0.55 (−1.18–0.09)	0	0.09	
Duration					
≥8 weeks	4	−1.54 (−2.50 to −0.59)	79	0.002	0.03
<8 weeks	5	−0.40 (−0.80–0.01)	0	0.05	
SJ					
Age					
≥18	1	−0.82 (−1.74–0.10)	–	0.08	0.36
<18	5	−1.44 (−2.39 to −0.49)	80	0.003	

(Continued)

TABLE 5 | Continued

	Groups (n)	SMD (95% CI)	I ²	p	p _{diff}
Duration					
≥8 weeks	3	-2.01 (-3.24 to -0.78)	79	0.001	0.04
<8 weeks	3	-0.64 (-1.17 to -0.11)	0	0.02	
COD ≤3 turns					
Age					
≥18	1	0.66 (-0.35-1.68)	-	0.20	0.45
<18	5	1.19 (0.28-2.10)	80	0.01	
Playing level					
Professional	5	1.19 (0.28-2.10)	80	0.01	0.45
Amateur	1	0.66 (-0.35-1.68)	-	0.20	
Duration					
≥8 weeks	3	1.84 (1.02-2.66)	63	<0.001	0.002
<8 weeks	3	0.31 (-0.25-0.87)	0	0.28	
COD >3 turns					
Age					
≥18	2	0.82 (0.16-1.48)	10	0.01	0.13
<18	5	1.43 (1.00-1.86)	8	<0.001	
Playing level					
Professional	4	1.53 (1.12-1.95)	0	<0.001	0.02
Amateur	3	0.63 (0.01-1.24)	0	0.05	
Duration					
≥8 weeks	4	1.53 (1.12-1.95)	0	<0.001	0.02
<8 weeks	3	0.63 (0.01-1.24)	0	0.05	

CI, confidence interval; CMJ, countermovement jump; COD, change of direction; I², heterogeneity; p_{diff}, test for subgroup difference; p, test for overall effect; SJ, squat jump; SMD, standardized mean difference. The bold values represents statistical significance between subgroups.

jump (SMD = 2.01 vs. 0.64, $p = 0.04$), COD with ≤3 turns (SMD = 1.84 vs. 0.31, $p = 0.002$), and COD >3 turns (SMD = 1.53 vs. 0.63, $p = 0.02$) was noted after ≥8 weeks compared to <8 weeks of CT (Table 5).

Regarding the competitive level of soccer players, a greater improvement in 30-m linear sprint time (SMD = 1.58 vs. 0.08, $p = 0.03$) and COD >3 turns (SMD = 1.53 vs. 0.63, $p = 0.02$) was noted in professional compared to amateur players after CT (Table 5).

Adverse Effect of Complex Training Intervention

Among the included studies, no injuries related to CT were reported. One study (Spinetti et al., 2019) reported the total RPE-load for CT and RT intervention. A total RPE-load of 25,201 was reported after 48 sessions of CT, while a total RPE-load of 25,364 was reported after RT.

DISCUSSION

Our meta-analyses indicated that compared to soccer training alone, CT induced moderate to large improvements in linear sprinting time, vertical jump height and sprinting with COD time in soccer players. The potentially beneficial effects derived

from CT seem particularly applicable to younger players (<18 vs. ≥18 years, for linear sprinting), after longer training interventions (<8 vs. ≥8 weeks, for jumping and COD) and among players with greater competitive level (professional vs. amateur, for linear sprinting and with COD). Further, among the included studies, no injuries related to CT were reported. Overall, CT seems safe and effective compared to soccer training alone for the improvement of the physical fitness of soccer players.

The improvements in linear sprinting, vertical jumping and sprinting with COD performance following CT may be related to adaptative mechanisms similar to those induced by RT and PT alone, including maximal strength, hormonal milieu, structural, and neuromechanical adaptations, which may be potentiated through a cumulative post-activation performance enhancement effect induced with CT (Sale, 2002; Robbins, 2005; Carter and Greenwood, 2014). Indeed, several CT interventions included in this systematic review observed significant IRM improvements in soccer players to a similar extent as RT (Faude et al., 2013; Brito et al., 2014; Spinetti et al., 2016; Hammami et al., 2017a, 2019; Chatzinikolaou et al., 2018). As maximal strength is strongly related to sprint, jump, and COD in male soccer players (Wisløff et al., 2004), improvements in maximal strength may be an additional and relevant benefit derived from CT. In addition, improvements in sprint, jump,

and COD performance after CT may be related to hormonal (Beaven et al., 2011; Ali et al., 2019) (e.g., testosterone increase) and cellular adaptations favorable to strength-power generation. For example, type IIx muscle fibers (i.e., those with greater contraction velocity, power, and rate of force development compared to type IIA and type I fibers) (Bottinelli et al., 1996; Harridge et al., 1996) may be favorably affected (i.e., greater preservation) by CT, even when compared to RT (Stasinaki et al., 2015). Indeed, an early study (Adams et al., 1993) reported that 19 weeks of heavy RT reduced percentage of type IIx fibers, whereas a more recent study (Stasinaki et al., 2015) reported a preservation of type IIx fibers after CT, similar to other studies that have included PT or CT (Macaluso et al., 2012, 2014; Grgic et al., 2020). Although not analyzed in the current study, the combination of RT and PT during CT may allow the summation of cellular and structural adaptations induced by RT and PT. Indeed, increased leg muscle volume was observed in soccer players after CT (Hammami et al., 2017a,b; Hammami et al., 2019). Although increases in muscle volume are usually achieved after RT interventions (ACSM, 2009; Krzysztosfik et al., 2019), recent research suggest that PT may also induce significant hypertrophic effects (Grgic et al., 2020; Gumpenberger et al., 2020). Further, such structural adaptations after RT and PT seem not to interfere with neuromechanical adaptations (ACSM, 2009; Markovic and Mikulic, 2010). In fact, CT may favor energy transfer between concentric and eccentric muscle actions, providing better coordination and synchronization of active muscle groups to improve and enhance motor skills (Cronin et al., 2001; Robbins, 2005). Such improvements may include sprinting, vertical jumping and sprinting with COD performance, in addition to other motor skills enhancements such as kicking ability (Cavaco et al., 2014; García-Pinillos et al., 2014). Altogether, CT seems an effective training strategy for the enhancement of sprinting, jumping and sprint with COD. However, future studies should elucidate the effectiveness of CT on such outcomes among soccer players, particularly when compared to other training methods, including RT.

A cumulative post-activation performance enhancement effect induced with CT (Sale, 2002; Robbins, 2005) may help explain the performance enhancement (Tillin and Bishop, 2009; Petisco et al., 2019). Further, as CT combine higher (i.e., RT) and lower loads (i.e., PT) this may have optimized the force-velocity curve of soccer players (Cormie et al., 2011) by ensuring that the prescribed training addressed the two broad components of the continuum. Considering the relevance of the force-velocity spectrum parameters among soccer players (Jimenez-Reyes et al., 2020), it seems plausible that an optimization of the force-velocity spectrum may help to explain the improvements in performance after CT (Jiménez-Reyes et al., 2017; Jimenez-Reyes et al., 2019). Such a combination of high-load low-velocity and low-load high-velocity exercises during CT may favor recruitment of fast-twitch muscle fibers (Gołaś et al., 2016), which are particularly important during maximal-intensity and short-duration actions (e.g., vertical jumping) (Fry et al., 2003; Macaluso et al., 2012). In contrast to our findings, a previous meta-analysis (Bauer et al., 2019) did not observe jumping improvements after CT. However, Bauer et al. (2019) included studies involving a myriad of

participants (e.g., schoolchildren, recreational trained students, untrained women, athletes from different sports) and compared the effects of CT with mixed training methods (e.g., RT, PT, mixed methods). In this sense, the results from our meta-analysis may offer a finding specific for soccer players (i.e., male soccer players), a finding that could have been potentially distorted amongst a wider population of athletes or the general population. On the other hand, extrapolation of current findings to other populations should proceed with caution.

Subgroup analyses were conducted to determine the effects of CT on physical fitness according to soccer player's chronological age, competitive level and duration of training programmes. Regarding the subgroup analysis according to the soccer players chronological age (<18 vs. ≥18 years), greater performance enhancements were noted after CT among players <18 years of age. The greater improvement among younger players may be due to the fact that players ≥18 years were already in the later stage of physical development and, hence, possessed a lower ceiling for further adaptation. In addition, players >18 years of age were likely to have performed RT and PT for a longer period of time (i.e., greater training age), experiencing diminishing return from CT (Carter and Greenwood, 2014) in the process. However, our subgroup analysis showed that professional players achieved greater performance improvements after CT compared to amateur players. Professional athletes have been shown to be better responders to post-activation performance enhancement (Carter and Greenwood, 2014). Indeed, professional athletes also tend to possess greater strength levels than amateur athletes, which might have enabled them to maximize the benefits of post-activation performance enhancement with CT (Seitz et al., 2014; Seitz and Haff, 2016; Suchomel et al., 2016). In addition, professional players' may possess a greater proportion of fast-twitch muscle fibers compared to amateur soccer players (Ostojic, 2004), rendering greater benefits from post-activation performance enhancement with CT (Seitz et al., 2014; Seitz and Haff, 2016; Suchomel et al., 2016). Furthermore, professional players may train with greater intensity, with professional athletes possessing greater intent to lift ballistically compared to non-professional athletes (Suchomel et al., 2018). In addition, CT was performed concurrently to soccer practices. It is possible that the number of soccer training sessions and games were greater in professional compared to non-professional players, with the former performing also greater number of sprints, jumps, and high-intensity actions. This may have also concurred to greater overall better neuromuscular adaptations in the professional vs. amateur athletes. Finally, regarding the subgroup analysis according to CT programme duration (<8 vs. ≥8 weeks), greater physical fitness improvements were noted after longer training interventions, in line with previous literature (Aagaard et al., 2001; Blazevich et al., 2007; Bolger et al., 2015). Although not surprising, current findings confirm that longer CT interventions may also induce larger gains in the physical fitness of male soccer players, particularly for jumping and sprints with COD.

Some limitations are acknowledged. Firstly, the findings of our systematic review suggest the absence of CT studies involving female soccer players. Therefore, current findings

should not be simply extrapolated to female athletes. Future CT research should involve female soccer players to overcome this shortcoming in literature. Another major shortcoming noted in this systematic review is the lack of studies comparing CT to RT. Future research should focus on this comparison.

CONCLUSION

The findings of this meta-analysis suggests that supplementing the regular soccer training sessions with CT improves athlete's physical fitness (i.e., sprinting, jumping, and COD ability) compared to soccer training alone. A longer duration of CT (≥ 8 weeks) seems to be optimal in improving the physical abilities of soccer players. Professional players and <18 years players may benefit more from CT program.

REFERENCES

- Aagaard, P., Andersen, J. L., Dyhre-Poulsen, P., Leffers, A. M., Wagner, A., Magnusson, S. P., et al. (2001). A mechanism for increased contractile strength of human pennate muscle in response to strength training: changes in muscle architecture. *J. Physiol.* 534, 613–623. doi: 10.1111/j.1469-7793.2001.t01-1-00613.x
- ACSM (2009). American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med. Sci. Sports Exerc.* 41, 687–708. doi: 10.1249/MSS.0b013e3181915670
- Adams, G. R., Hather, B. M., Baldwin, K. M., and Dudley, G. A. (1993). Skeletal muscle myosin heavy chain composition and resistance training. *J. Appl. Physiol.* 74, 911–915. doi: 10.1152/jappl.1993.74.2.911
- Ali, K., Verma, S., Ahmad, I., Singla, D., Saleem, M., and Hussain, M. E. (2019). Comparison of complex versus contrast training on steroid hormones and sports performance in male soccer players. *J. Chiropr. Med.* 18, 131–138. doi: 10.1016/j.jcm.2018.12.001
- Arnason, A., Sigurdsson, S. B., Gudmundsson, A., Holme, I., Engebretsen, L., and Bahr, R. (2004). Physical fitness, injuries, and team performance in soccer. *Med. Sci. Sports Exerc.* 36, 278–285. doi: 10.1249/01.MSS.0000113478.92945.CA
- Asadi, A., Arazi, H., Young, W. B., and De Villarreal, E. S. (2016). The effects of plyometric training on change-of-direction ability: a meta-analysis. *Int. J. Sports Physiol. Perf.* 11, 563–573. doi: 10.1123/ijsp.2015-0694
- Barnes, C., Archer, D. T., Hogg, B., Bush, M., and Bradley, P. S. (2014). The evolution of physical and technical performance parameters in the English Premier League. *Int. J. Sports Med.* 35, 1095–1100. doi: 10.1055/s-0034-1375695
- Bauer, P., Uebellacker, F., Mitter, B., Aigner, A. J., Hasenoehrl, T., Ristl, R., et al. (2019). Combining higher-load and lower-load resistance training exercises: a systematic review and meta-analysis of findings from complex training studies. *J. Sci. Med. Sport* 22, 838–851. doi: 10.1016/j.jsams.2019.01.006
- Beaven, C. M., Gill, N. D., Ingram, J. R., and Hopkins, W. G. (2011). Acute salivary hormone responses to complex exercise bouts. *J. Strength Cond. Res.* 25, 1072–1078. doi: 10.1519/JSC.0b013e3181bf4414
- Blazevich, A. J., Cannavan, D., Coleman, D. R., and Horne, S. (2007). Influence of concentric and eccentric resistance training on architectural adaptation in human quadriceps muscles. *J. Appl. Physiol.* 103, 1565–1575. doi: 10.1152/japplphysiol.00578.2007
- Bolger, R., Lyons, M., Harrison, A. J., and Kenny, I. C. (2015). Sprinting performance and resistance-based training interventions: a systematic review. *J. Strength Cond. Res.* 29, 1146–1156. doi: 10.1519/JSC.0000000000000720
- Borenstein, M., Hedges, L. V., Higgins, J. P., and Rothstein, H. R. (2009). “Identifying and Quantifying Heterogeneity,” in *Introduction to Meta-Analysis*, eds M. Borenstein, L. V. Hedges, J. P. Higgins and H. R. Rothstein (Chichester, UK: John Wiley & Sons), 107–125.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author.

AUTHOR CONTRIBUTIONS

All authors made significant contributions, including preparation of the first draft of the manuscript, data collection, analysis of data, interpretation of data, and/or provided meaningful revision and feedback, read, and approved the final manuscript.

ACKNOWLEDGMENTS

The authors want to thank Gopal Kumar for providing help in initial extraction of studies for this meta-analysis.

- Bottinelli, R., Canepari, M., Pellegrino, M. A., and Reggiani, C. (1996). Force-velocity properties of human skeletal muscle fibres: myosin heavy chain isoform and temperature dependence. *J. Physiol.* 495(Pt 2), 573–586. doi: 10.1113/jphysiol.1996.sp021617
- Brito, J., Vasconcellos, F., Oliveira, J., Krustup, P., and Rebelo, A. (2014). Short-term performance effects of three different low-volume strength-training programmes in college male soccer players. *J. Hum. Kinet.* 40, 121–128. doi: 10.2478/hukin-2014-0014
- Carter, J., and Greenwood, M. (2014). Complex training reexamined: review and recommendations to improve strength and power. *Strength Cond. J.* 36, 11–19. doi: 10.1519/SSC.0000000000000036
- Castagna, C., D'ottavio, S., and Abt, G. (2003). Activity profile of young soccer players during actual match play. *J. Strength Cond. Res.* 17, 775–780. doi: 10.1519/00124278-200311000-00024
- Cavaco, B., Sousa, N., Dos Reis, V. M., Garrido, N., Saavedra, F., Mendes, R., et al. (2014). Short-term effects of complex training on agility with the ball, speed, efficiency of crossing and shooting in youth soccer players. *J. Hum. Kinet.* 43, 105–112. doi: 10.2478/hukin-2014-0095
- Chatziniolaou, A., Michaloglou, K., Avloniti, A., Leontini, D., Deli, C. K., Vlachopoulos, D., et al. (2018). The trainability of adolescent soccer players to brief periodized complex training. *Int. J. Sports Physiol. Perform.* 13, 645–655. doi: 10.1123/ijsp.2017-0763
- Cohen, J. (1988). “The t tests for means,” in *Statistical Power Analysis for the Behavioral Sciences, 2nd Edn.*, ed J. Cohen (New York, NY: Routledge), 19–40.
- Cormie, P., McGuigan, M. R., and Newton, R. U. (2011). Developing maximal neuromuscular power: part 1—biological basis of maximal power production. *Sports Med.* 41, 17–38. doi: 10.2165/11537690-000000000-00000
- Cormier, P., Freitas, T. T., Rubio-Arias, J. A., and Alcaraz, P. E. (2020). Complex and contrast training: does strength and power training sequence affect performance-based adaptations in team sports? a systematic review and meta-analysis. *J. Strength Cond. Res.* 34, 1461–1479. doi: 10.1519/JSC.0000000000003493
- Cronin, J., Mcnair, P. J., and Marshall, R. N. (2001). Velocity specificity, combination training and sport specific tasks. *J. Sci. Med. Sport* 4, 168–178. doi: 10.1016/S1440-2440(01)80027-X
- De Hoyo, M., Gonzalo-Skok, O., Sañudo, B., Carrascal, C., Plaza-Armas, J. R., Camacho-Candil, F., et al. (2016). Comparative effects of in-season full-back squat, resisted sprint training, and plyometric training on explosive performance in u-19 elite soccer players. *J. Strength Cond. Res.* 30, 368–377. doi: 10.1519/JSC.0000000000001094
- De Villarreal, E. S., Requena, B., and Cronin, J. B. (2012). The effects of plyometric training on sprint performance: a meta-analysis. *J. Strength Cond. Res.* 26, 575–584. doi: 10.1519/JSC.0b013e318220fd03

- Deeks, J. J., Higgins, J. P., and Altman, D. G. (2008). "Analysing data and undertaking meta-analyses," in *Cochrane Handbook for Systematic Reviews of Interventions*, eds J. P. Higgins and S. Green (Chichester, UK: John Wiley & Sons), 243–296.
- Docherty, D., Robbins, D., and Hodgson, M. (2004). Complex training revisited: a review of its current status as a viable training approach. *Strength Cond. J.* 26, 52–57. doi: 10.1519/00126548-200412000-00011
- Drevon, D., Fursa, S. R., and Malcolm, A. L. (2017). Intercoder reliability and validity of webplotdigitizer in extracting graphed data. *Behav. Modif.* 41, 323–339. doi: 10.1177/0145445516673998
- Faude, O., Koch, T., and Meyer, T. (2012). Straight sprinting is the most frequent action in goal situations in professional football. *J. Sports Sci.* 30, 625–631. doi: 10.1080/02640414.2012.665940
- Faude, O., Roth, R., Di Giovine, D., Zahner, L., and Donath, L. (2013). Combined strength and power training in high-level amateur football during the competitive season: a randomised-controlled trial. *J. Sports Sci.* 31, 1460–1467. doi: 10.1080/02640414.2013.796065
- Fleck, S., and Kontor, K. (1986). Complex training. *Strength Cond. J.* 8, 66–72. doi: 10.1519/0744-0049(1986)008<0066:CT>2.3.CO;2
- Fleck, S. J., and Kraemer, W. J. (2004). *Designing Resistance Training Programs*. Champaign, IL: Human Kinetics.
- Freitas, T. T., Martínez-Rodríguez, A., Calleja-González, J., and Alcaraz, P. E. (2017). Short-term adaptations following complex training in team-sports: a meta-analysis. *PLoS ONE* 12:e0180223. doi: 10.1371/journal.pone.0180223
- Fry, A. C., Schilling, B. K., Staron, R. S., Hagerman, F. C., Hikida, R. S., and Thrush, J. T. (2003). Muscle fiber characteristics and performance correlates of male Olympic-style weightlifters. *J. Strength Cond. Res.* 17, 746–754. doi: 10.1519/00124278-200311000-00020
- García-Pinillos, F., Martínez-Amat, A., Hita-Contreras, F., Martínez-López, E. J., and Latorre-Román, P. A. (2014). Effects of a contrast training program without external load on vertical jump, kicking speed, sprint, and agility of young soccer players. *J. Strength Cond. Res.* 28, 2452–2460. doi: 10.1519/JSC.0000000000000452
- Golaś, A., Maszczyk, A., Zajac, A., Mikołajec, K., and Stastny, P. (2016). Optimizing post activation potentiation for explosive activities in competitive sports. *J. Hum. Kinet.* 52, 95–106. doi: 10.1515/hukin-2015-0197
- Grgic, J., Schoenfeld, B. J., and Mikulic, P. (2020). Effects of plyometric vs. resistance training on skeletal muscle hypertrophy: a review. *J. Sport Health Sci.* doi: 10.1016/j.jshs.2020.06.010. [Epub ahead of print].
- Gumpenberger, M., Wessner, B., Graf, A., Narici, M. V., Fink, C., Braun, S., et al. (2020). Remodeling the skeletal muscle extracellular matrix in older age-effects of acute exercise stimuli on gene expression. *Int. J. Mol. Sci.* 21:7089. doi: 10.3390/ijms21197089
- Hammami, M., Gaamouri, N., Shephard, R. J., and Chelly, M. S. (2019). Effects of contrast strength vs. plyometric training on lower-limb explosive performance, ability to change direction and neuromuscular adaptation in soccer players. *J. Strength Cond. Res.* 33, 2094–2103. doi: 10.1519/JSC.00000000000002425
- Hammami, M., Negra, Y., Shephard, R. J., and Chelly, M. S. (2017a). The effect of standard strength vs. contrast strength training on the development of sprint, agility, repeated change of direction, and jump in junior male soccer players. *J. Strength Cond. Res.* 31, 901–912. doi: 10.1519/JSC.00000000000001815
- Hammami, M., Negra, Y., Shephard, R. J., and Chelly, M. S. (2017b). Effects of leg contrast strength training on sprint, agility and repeated change of direction performance in male soccer players. *J. Sports Med. Phys. Fitness* 57, 1424–1431. doi: 10.23736/S0022-4707.17.06951-1
- Harridge, S. D., Bottinelli, R., Canepari, M., Pellegrino, M. A., Reggiani, C., Esbjörnsson, M., et al. (1996). Whole-muscle and single-fibre contractile properties and myosin heavy chain isoforms in humans. *Pflugers Arch.* 432, 913–920. doi: 10.1007/s004240050215
- Healy, R., and Comyns, T. M. (2017). The application of postactivation potentiation methods to improve sprint speed. *Strength Cond. J.* 39, 1–9. doi: 10.1519/SSC.0000000000000276
- Higgins, J. P. (2008). Commentary: heterogeneity in meta-analysis should be expected and appropriately quantified. *Int. J. Epidemiol.* 37, 1158–1160. doi: 10.1093/ije/dyn204
- Higgins, J. P., Deeks, J. J., and Altman, D. G. (2008). "Special Topics in Statistics," in *Cochrane Handbook for Systematic Reviews of Interventions*, eds J. P. Higgins and S. Green (Chichester, UK: John Wiley & Sons), 481–529. doi: 10.1002/9780470712184.ch16
- Higgins, J. P., Thompson, S. G., Deeks, J. J., and Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *BMJ* 327, 557–560. doi: 10.1136/bmj.327.7414.557
- Hodgson, M., Docherty, D., and Robbins, D. (2005). Post-activation potentiation: underlying physiology and implications for motor performance. *Sports Med.* 35, 585–595. doi: 10.2165/00007256-200535070-00004
- Hopkins, W. G., Marshall, S. W., Batterham, A. M., and Hanin, J. (2009). Progressive statistics for studies in sports medicine and exercise science. *Med. Sci. Sports Exerc.* 41, 3–13. doi: 10.1249/MSS.0b013e31818cb278
- Izquierdo, M., Häkkinen, K., Gonzalez-Badillo, J. J., Ibáñez, J., and Gorostiaga, E. M. (2002). Effects of long-term training specificity on maximal strength and power of the upper and lower extremities in athletes from different sports. *Eur. J. Appl. Physiol.* 87, 264–271. doi: 10.1007/s00421-002-0628-y
- Jimenez-Reyes, P., García-Ramos, A., Parraga-Montilla, J. A., Morcillo-Losa, J. A., Cuadrado-Penañiel, V., Castano-Zambudio, A., et al. (2020). Seasonal changes in the sprint acceleration force-velocity profile of elite male soccer players. *J. Strength Cond. Res.* doi: 10.1519/JSC.0000000000003513. [Epub ahead of print].
- Jiménez-Reyes, P., Samozino, P., Brughelli, M., and Morin, J. B. (2017). Effectiveness of an individualized training based on force-velocity profiling during jumping. *Front. Physiol.* 7:677. doi: 10.3389/fphys.2016.00677
- Jimenez-Reyes, P., Samozino, P., and Morin, J. B. (2019). Optimized training for jumping performance using the force-velocity imbalance: individual adaptation kinetics. *PLoS ONE* 14:e0216681. doi: 10.1371/journal.pone.0216681
- Kobal, R., Loturco, I., Barroso, R., Gil, S., Cuniyochi, R. R., Ugrinowitsch, C., et al. (2017). Effects of different combinations of strength, power, and plyometric training on the physical performance of elite young soccer players. *J. Strength Cond. Res.* 31, 1468–1476. doi: 10.1519/JSC.0000000000001609
- Kontopantelis, E., Springate, D. A., and Reeves, D. (2013). A re-analysis of the Cochrane Library data: the dangers of unobserved heterogeneity in meta-analyses. *PLoS ONE* 8:e69930. doi: 10.1371/journal.pone.0069930
- Krzysztofik, M., Wilk, M., Wojdała, G., and Golaś, A. (2019). Maximizing muscle hypertrophy: a systematic review of advanced resistance training techniques and methods. *Int. J. Environ. Res. Public Health* 16:4897. doi: 10.3390/ijerph16244897
- Liberati, A., Altman, D. G., Tetzlaff, J., Mulrow, C., Gøtzsche, P. C., Ioannidis, J. P., et al. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J. Clin. Epidemiol.* 62, e1–34. doi: 10.1016/j.jclinepi.2009.06.006
- Macaluso, F., Isaacs, A. W., Di Felice, V., and Myburgh, K. H. (2014). Acute change of titin at mid-sarcomere remains despite 8 wk of plyometric training. *J. Appl. Physiol.* 116, 1512–1519. doi: 10.1152/jappphysiol.00420.2013
- Macaluso, F., Isaacs, A. W., and Myburgh, K. H. (2012). Preferential type II muscle fiber damage from plyometric exercise. *J. Athl. Train* 47, 414–420. doi: 10.4085/1062-6050-47.4.13
- Maio Alves, J. M., Rebelo, A. N., Abrantes, C., and Sampaio, J. (2010). Short-term effects of complex and contrast training in soccer players' vertical jump, sprint, and agility abilities. *J. Strength Cond. Res.* 24, 936–941. doi: 10.1519/JSC.0b013e3181c7c5fd
- Marios, C., Smilios, I., Sotiropoulos, K., Volaklis, K., Theophilos, P., and Tokmakidis, S. (2006). The effects of resistance training on the physical capacities of adolescent soccer players. *J. Strength Cond. Res.* 20, 783–791. doi: 10.1519/00124278-200611000-00010
- Markovic, G., and Mikulic, P. (2010). Neuro-musculoskeletal and performance adaptations to lower-extremity plyometric training. *Sports Med.* 40, 859–895. doi: 10.2165/11318370-000000000-00000
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., et al. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst. Rev.* 4:1. doi: 10.1186/2046-4053-4-1
- Ostojic, S. M. (2004). Elite and nonelite soccer players: pre-seasonal physical and physiological characteristics. *Res. Sports Med.* 12, 143–150. doi: 10.1080/15438620490460495

- Pagaduan, J., and Pojskic, H. (2020). A meta-analysis on the effect of complex training on vertical jump performance. *J. Hum. Kinet.* 71, 255–265. doi: 10.2478/hukin-2019-0087
- Petisco, C., Ramirez-Campillo, R., Hernández, D., Gonzalo-Skok, O., Nakamura, F. Y., and Sanchez-Sanchez, J. (2019). Post-activation potentiation: effects of different conditioning intensities on measures of physical fitness in male young professional soccer players. *Front. Psychol.* 10:1167. doi: 10.3389/fpsyg.2019.01167
- Prieske, O., Behrens, M., Chaabene, H., Granacher, U., and Maffiuletti, N. A. (2020). Time to differentiate postactivation “potentiation” from “performance enhancement” in the strength and conditioning community. *Sports Med.* 50, 1559–1565. doi: 10.1007/s40279-020-01300-0
- Ramirez-Campillo, R., Castillo, D., Raya-González, J., Moran, J., De Villarreal, E. S., and Lloyd, R. S. (2020a). Effects of plyometric jump training on jump and sprint performance in young male soccer players: a systematic review and meta-analysis. *Sports Med.* 50, 2125–2143. doi: 10.1007/s40279-020-01337-1
- Ramirez-Campillo, R., Sanchez-Sanchez, J., Romero-Moraleda, B., Yanci, J., García-Hermoso, A., and Manuel Clemente, F. (2020b). Effects of plyometric jump training in female soccer player’s vertical jump height: a systematic review with meta-analysis. *J. Sports Sci.* 38, 1475–1487. doi: 10.1080/02640414.2020.1745503
- Robbins, D. W. (2005). Postactivation potentiation and its practical applicability: a brief review. *J. Strength Cond. Res.* 19, 453–458. doi: 10.1519/R-14653.1
- Sale, D. G. (2002). Postactivation potentiation: role in human performance. *Exerc. Sport Sci. Rev.* 30, 138–143. doi: 10.1097/00003677-200207000-00008
- Sander, A., Keiner, M., Wirth, K., and Schmidtbleicher, D. (2013). Influence of a 2-year strength training programme on power performance in elite youth soccer players. *Eur. J. Sport Sci.* 13, 445–451. doi: 10.1080/17461391.2012.742572
- Seitz, L. B., De Villarreal, E. S., and Haff, G. G. (2014). The temporal profile of postactivation potentiation is related to strength level. *J. Strength Cond. Res.* 28, 706–715. doi: 10.1519/JSC.0b013e3182a73ea3
- Seitz, L. B., and Haff, G. G. (2016). Factors modulating post-activation potentiation of jump, sprint, throw, and upper-body ballistic performances: a systematic review with meta-analysis. *Sports Med.* 46, 231–240. doi: 10.1007/s40279-015-0415-7
- Silva, J., Nassis, G., and Rebelo, A. (2015). Strength training in soccer with a specific focus on highly trained players. *Sports Med. Open* 1:17. doi: 10.1186/s40798-015-0006-z
- Smart, N. A., Waldron, M., Ismail, H., Giallauria, F., Vigorito, C., Cornelissen, V., et al. (2015). Validation of a new tool for the assessment of study quality and reporting in exercise training studies: TESTEX. *Int. J. Evid. Based Healthc.* 13, 9–18. doi: 10.1097/XEB.000000000000020
- Spinetti, J., Figueiredo, T., Bastos, D. E. O. V., Assis, M., Fernandes, D. E. O. L., Miranda, H., et al. (2016). Comparison between traditional strength training and complex contrast training on repeated sprint ability and muscle architecture in elite soccer players. *J. Sports Med. Phys. Fitness* 56, 1269–1278.
- Spinetti, J., Figueiredo, T., Willardson, J., Bastos De Oliveira, V., Assis, M., Fernandes De Oliveira, L., et al. (2019). Comparison between traditional strength training and complex contrast training on soccer players. *J. Sports Med. Phys. Fitness* 59, 42–49. doi: 10.23736/S0022-4707.18.07934-3
- Stasinaki, A. N., Gloumis, G., Spengos, K., Blazevich, A. J., Zaras, N., Georgiadis, G., et al. (2015). Muscle strength, power, and morphologic adaptations after 6 weeks of compound vs. complex training in healthy men. *J. Strength Cond. Res.* 29, 2559–2569. doi: 10.1519/JSC.0000000000000917
- Stølen, T., Chamari, K., Castagna, C., and Wisløff, U. (2005). Physiology of soccer: an update. *Sports Med.* 35, 501–536. doi: 10.2165/00007256-200535060-00004
- Suchomel, T. J., Nimphius, S., Bellon, C. R., and Stone, M. H. (2018). The importance of muscular strength: training considerations. *Sports Med.* 48, 765–785. doi: 10.1007/s40279-018-0862-z
- Suchomel, T. J., Nimphius, S., and Stone, M. H. (2016). The importance of muscular strength in athletic performance. *Sports Med.* 46, 1419–1449. doi: 10.1007/s40279-016-0486-0
- Taylor, J. B., Ford, K. R., Schmitz, R. J., Ross, S. E., Ackerman, T. A., and Shultz, S. J. (2018). Sport-specific biomechanical responses to an ACL injury prevention programme: a randomised controlled trial. *J. Sports Sci.* 36, 2492–2501. doi: 10.1080/02640414.2018.1465723
- Tillin, N. A., and Bishop, D. (2009). Factors modulating post-activation potentiation and its effect on performance of subsequent explosive activities. *Sports Med.* 39, 147–166. doi: 10.2165/00007256-200939020-00004
- Van De Hoef, P. A., Brauers, J. J., Van Smeden, M., Backx, F. J. G., and Brink, M. S. (2019). The effects of lower-extremity plyometric training on soccer-specific outcomes in adult male soccer players: a systematic review and meta-analysis. *Int. J. Sports Physiol. Perform.* 15, 1–15. doi: 10.1123/ijspp.2019-0565
- Vlachopoulos, D., Barker, A. R., Ubago-Guisado, E., Williams, C. A., and Gracia-Marco, L. (2018). The effect of a high-impact jumping intervention on bone mass, bone stiffness and fitness parameters in adolescent athletes. *Arch. Osteoporosis* 13:128. doi: 10.1007/s11657-018-0543-4
- Wisløff, U., Castagna, C., Helgerud, J., Jones, R., and Hoff, J. (2004). Strong correlation of maximal squat strength with sprint performance and vertical jump height in elite soccer players. *Br. J. Sports Med.* 38, 285–288. doi: 10.1136/bjism.2002.002071

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Thapa, Lum, Moran and Ramirez-Campillo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.